

BEVERLY HILLS PEDIATRICS

DOSING CHART FOR **TYLENOL** (to be given every 4 hours as needed)

Age	weight	infant drops 80mg/0.8ml	childrens liquid susp 160mg/5ml	chewable tabs 80mg
0-3 mo	6-11 lbs	0.4ml	1/4 tsp	--
4-9 mo	12-15 lbs	0.8ml	1/2 tsp	one tab
9-12 mo	16-21lbs	1.2 ml	3/4 tsp	1.5 tabs
12-23 mo	22-32lbs	1.6ml	one tsp	2 tabs
2-3 yrs	33-40lbs		1 1/2 tsp	3 tabs
4-5yrs	41-54lbs		2 tsp	4 tabs
6-8 yrs	55-64 lbs		2 1/2 tsp	5 tabs
9-10yrs	65-88lbs		3 tsp	6 tabs

DOSING CHART FOR **ADVIL and MOTRIN** (to be given every 6 hours as needed)

****do not give to children under 6months****

Age	weight	infant drops 50mg/1.25ml	childrens liquid susp 100mg/5ml	chew tabs 50mg	jr chewable tab 100mg
6-9 mo	12-15 lbs	1.25ml	1/2 tsp	one tab	
9-12 mo	16-21lbs	1.875 ml	3/4 tsp		
12-23 mo	22-32lbs	2.5ml	one tsp	2 tabs	one tab
2-3 yrs	33-40lbs		1 1/2 tsp	3 tabs	1 1/2 tab
4-5yrs	41-54lbs		2 tsp	4 tabs	2 tabs
6-8 yrs	55-64 lbs		2 1/2 tsp	5 tabs	2 1/2 tabs
9-10yrs	65-88lbs		3 tsp	6 tabs	3 tabs

DOSING CHART FOR **CHILDREN'S LIQUID PREPARATIONS** (cough, cold etc.)

(Benadryl, Dimetapp, Triaminic etc)

****COUGH AND COLD MEDICATIONS ARE NOT RECOMMENDED FOR CHILDREN UNDER 2 YEARS OF AGE****

****Benadryl may be used in children under 2 years of age for allergy purposes only****

Age	weight	dose	frequency
6-11 mo	12-17 lbs	1/2 tsp	every 6 hrs
1-2 yrs	18-30 lbs	one tsp	every 6 hrs
3-6 yrs	31-44 lbs	1 1/2 tsp	every 6 hrs
6-12 yrs	45 lbs and up	2 tsp	every 6 hrs